

## Relations between Distorted and Original Angles in STR

Florentin Smarandache

Arts and Science Division, University of New Mexico, 705 Gurley Ave., Gallup, NM 87301, USA. E-mail: smarand@unm.edu

Using the Oblique-Length Contraction Factor, which is a generalization of Lorentz Contraction Factor, one shows several trigonometric relations between distorted and original angles of a moving object lengths in the Special Theory of Relativity.

### 1 Introduction

The lengths at oblique angle to the motion are contracted with the Oblique-Length Contraction Factor  $OC(v, \theta)$ , defined as [1-2]:

$$OC(v, \theta) = \sqrt{C(v)^2 \cos^2 \theta + \sin^2 \theta} \quad (1)$$

where  $C(v)$  is just Lorentz Factor:

$$C(v) = \sqrt{1 - \frac{v^2}{c^2}} \in [0, 1] \text{ for } v \in [0, c]. \quad (2)$$

Of course

$$0 \leq OC(v, \theta) \leq 1. \quad (3)$$

The Oblique-Length Contraction Factor is a generalization of Lorentz Contractor  $C(v)$ , because: when  $\theta = 0$ , or the length is moving along the motion direction, then  $OC(v, 0) = C(v)$ . Similarly

$$OC(v, \pi) = OC(v, 2\pi) = C(v). \quad (4)$$

Also, if  $\theta = \pi/2$ , or the length is perpendicular on the motion direction, then  $OC(v, \pi/2) = 1$ , i.e. no contraction occurs. Similarly  $OC(v, \frac{3\pi}{2}) = 1$ .

### 2 Tangential relations between distorted acute angles vs. original acute angles of a right triangle

Let's consider a right triangle with one of its legs along the motion direction (Fig. 1).




Fig. 1:

$$\tan \theta = \frac{\beta}{\gamma} \quad (5)$$

$$\tan(180^\circ - \theta) = -\tan \theta = -\frac{\beta}{\gamma} \quad (6)$$

After contraction of the side  $AB$  (and consequently contraction of the oblique side  $BC$ ) one gets (Fig. 2):




Fig. 2:

$$\tan(180^\circ - \theta') = -\tan \theta' = -\frac{\beta'}{\gamma'} = -\frac{\beta}{\gamma C(v)}. \quad (7)$$

Then:

$$\frac{\tan(180^\circ - \theta')}{\tan(180^\circ - \theta)} = \frac{-\frac{\beta}{\gamma C(v)}}{-\frac{\beta}{\gamma}} = \frac{1}{C(v)}. \quad (8)$$

Therefore

$$\tan(\pi - \theta') = -\frac{\tan(\pi - \theta)}{C(v)} \quad (9)$$

and consequently

$$\tan(\theta') = \frac{\tan(\theta)}{C(v)} \quad (10)$$

or

$$\tan(B') = \frac{\tan(B)}{C(v)} \quad (11)$$

which is the Angle Distortion Equation, where  $\theta$  is the angle formed by a side travelling along the motion direction and another side which is oblique on the motion direction.

The angle  $\theta$  is increased (i.e.  $\theta' > \theta$ ).

$$\tan \varphi = \frac{\gamma}{\beta} \quad \text{and} \quad \tan \varphi' = \frac{\gamma'}{\beta'} = \frac{\gamma C(v)}{\beta} \quad (12)$$

whence:

$$\frac{\tan \varphi'}{\tan \varphi} = \frac{\frac{\gamma C(v)}{\beta}}{\frac{\gamma}{\beta}} = C(v). \quad (13)$$

So we get the following Angle Distortion Equation:

$$\tan \varphi' = \tan \varphi \cdot C(v) \quad (14)$$

or

$$\tan C' = \tan C \cdot C(v) \quad (15)$$

where  $\varphi$  is the angle formed by one side which is perpendicular on the motion direction and the other one is oblique to the motion direction.

The angle  $\varphi$  is decreased (i.e.  $\varphi' < \varphi$ ). If the traveling right triangle is oriented the opposite way (Fig. 3)




Fig. 3:

$$\tan \theta = \frac{\beta}{\gamma} \quad \text{and} \quad \tan \varphi = \frac{\gamma}{\beta}. \quad (16)$$

Similarly, after contraction of side  $AB$  (and consequently contraction of the oblique side  $BC$ ) one gets (Fig. 4)

$$\tan \theta' = \frac{\beta'}{\gamma'} = \frac{\beta}{\gamma C(v)} \quad (17)$$

and

$$\tan \varphi' = \frac{\gamma'}{\beta'} = \frac{\gamma C(v)}{\beta} \quad (18)$$

$$\frac{\tan \theta'}{\tan \theta} = \frac{\frac{\beta}{\gamma C(v)}}{\frac{\beta}{\gamma}} = \frac{1}{C(v)} \quad (19)$$

or

$$\tan \theta' = \frac{\tan \theta}{C(v)} \quad (20)$$




Fig. 4:

and similarly

$$\frac{\tan \varphi'}{\tan \varphi} = \frac{\frac{\gamma C(v)}{\beta}}{\frac{\gamma}{\beta}} = C(v) \quad (21)$$

or

$$\tan \varphi' = \tan \varphi \cdot C(v). \quad (22)$$

Therefore one got the same Angle Distortion Equations for a right triangle traveling with one of its legs along the motion direction.

### 3 Tangential relations between distorted angles vs. original angles of a general triangle

Let's suppose a general triangle  $\Delta ABC$  is travelling at speed  $v$  along the side  $BC$  as in Fig. 5.




Fig. 5:

The height remains not contracted:  $AM \equiv A'M'$ . We can split this figure into two traveling right sub-triangles as in Fig. 6.

In the right triangles  $\Delta A'M'B'$  and respectively  $\Delta A'M'C'$  one has

$$\tan B' = \frac{\tan B}{C(v)} \quad \text{and} \quad \tan C' = \frac{\tan C}{C(v)}. \quad (23)$$

Also

$$\tan A'_1 = \tan A_1 C(v) \quad \text{and} \quad \tan A'_2 = \tan A_2 C(v). \quad (24)$$




Fig. 6:




Fig. 8:




Fig. 7:

But

$$\begin{aligned} \tan A' &= \tan(A'_1 + A'_2) = \frac{\tan A'_1 + \tan A'_2}{1 - \tan A'_1 \tan A'_2} \\ &= \frac{\tan A_1 C(\nu) + \tan A_2 C(\nu)}{1 - \tan A_1 C(\nu) \tan A_2 C(\nu)} \\ &= C(\nu) \cdot \frac{\tan A_1 + \tan A_2}{1 - \tan A_1 \tan A_2 C(\nu)^2} \\ &= C(\nu) \cdot \frac{\tan A_1 + \tan A_2}{1 - \tan A_1 \tan A_2} \cdot (1 - \tan A_1 \tan A_2) \\ &= C(\nu) \cdot \frac{\tan(A_1 + A_2)}{1} \cdot \frac{1 - \tan A_1 \tan A_2}{1 - \tan A_1 \tan A_2 C(\nu)^2}. \end{aligned}$$

$$\tan A' = C(\nu) \cdot \tan(A) \cdot \frac{1 - \tan A_1 \tan A_2}{1 - \tan A_1 \tan A_2 C(\nu)^2}. \quad (25)$$

We got

$$\tan A' = \tan(A) \cdot C(\nu) \cdot \frac{1 - \tan A_1 \tan A_2}{1 - \tan A_1 \tan A_2 C(\nu)^2} \quad (26)$$

Similarly we can split this Fig. 7 into two traveling right sub-triangles as in Fig. 8.

#### 4 Other relations between the distorted angles and the original angles

1. Another relation uses the Law of Sine in the triangles  $\Delta ABC$  and respectively  $\Delta A'B'C'$ :

$$\frac{\alpha}{\sin A} = \frac{\beta}{\sin B} = \frac{\gamma}{\sin C} \quad (27)$$

$$\frac{\alpha'}{\sin A'} = \frac{\beta'}{\sin B'} = \frac{\gamma'}{\sin C'}. \quad (28)$$

After substituting

$$\alpha' = \alpha C(\nu) \quad (29)$$

$$\beta' = \beta \theta C(\nu, C) \quad (30)$$

$$\gamma' = \gamma \theta C(\nu, B) \quad (31)$$

into the second relation one gets:

$$\frac{\alpha C(\nu)}{\sin A'} = \frac{\beta \theta C(\nu, C)}{\sin B'} = \frac{\gamma \theta C(\nu, B)}{\sin C'}. \quad (32)$$

Then we divide term by term the previous equalities:

$$\frac{\frac{\alpha}{\sin A}}{\frac{\alpha C(\nu)}{\sin A'}} = \frac{\frac{\beta}{\sin B}}{\frac{\beta \theta C(\nu, C)}{\sin B'}} = \frac{\frac{\gamma}{\sin C}}{\frac{\gamma \theta C(\nu, B)}{\sin C'}} \quad (33)$$

whence one has:

$$\begin{aligned} \frac{\sin A'}{\sin A \cdot C(\nu)} &= \frac{\sin B'}{\sin B \cdot \theta C(\nu, C)} \\ &= \frac{\sin C'}{\sin C \cdot \theta C(\nu, B)}. \end{aligned} \quad (34)$$

2. Another way:

$$A' = 180^\circ - (B' + C') \quad \text{and} \quad A = 180^\circ - (B + C) \quad (35)$$

$$\tan A' = \tan[180^\circ - (B' + C')] = -\tan(B' + C')$$

$$= -\frac{\tan B' + \tan C'}{1 - \tan B' \cdot \tan C'}$$

$$\begin{aligned}
&= -\frac{\frac{\tan B}{C(v)} + \frac{\tan C}{C(v)}}{1 - \tan B \cdot \tan C/C(v)^2} \\
&= -\frac{1}{C(v)} \cdot \frac{\tan B + \tan C}{1 - \tan B \cdot \tan C/C(v)^2} \\
&= -\frac{\tan(B+C)}{C(v)} \cdot \frac{1 - \tan B \tan C}{1 - \tan B \cdot \tan C/C(v)^2} \\
&= -\frac{-\tan[180^\circ - (B+C)]}{C(v)} \cdot \frac{1 - \tan B \cdot \tan C}{1 - \tan B \cdot \tan C/C(v)^2} \\
&= \frac{\tan A}{C(v)} \cdot \frac{1 - \tan B \cdot \tan C}{1 - \tan B \cdot \tan C/C(v)^2}.
\end{aligned}$$

We got

$$\tan A' = \frac{\tan A}{C(v)} \cdot \frac{1 - \tan B \cdot \tan C}{1 - \tan B \cdot \tan C/C(v)^2}. \quad (36)$$

Submitted on March 30, 2013 /Accepted on April 2, 2013

## References

1. Smarandache F. *New Relativistic Paradoxes and Open Questions*. Somipress, Fes, 1983.
2. Smarandache F. Oblique-Length Contraction Factor in the Special Theory of Relativity, *Progress in Physics*, 2013, v. 1, 60–62.
3. Einstein A. On the Electrodynamics of Moving Bodies. *Annalen der Physik*, 1905, v. 17, 891–921.
4. Smarandache F. *Absolute Theory of Relativity and Parameterized Special Theory of Relativity and Noninertial Multirelativity*. Somipress, Fes, 1982.