

Quantum Meteorological Resonator. Part II: Numerical Experiments

Mikhail L. Arushanov^{1,2}

¹Research Hydrometeorological Institute, 72 Chingiz Aitmatov Str., Tashkent 100052, Uzbekistan

²Central Asian University of Environmental and Climate Change Studies (Green University), Tashkent, Uzbekistan

E-mail: mikl-arushanov@rambler.ru

This paper represents a direct continuation of the author's fundamental theoretical framework (*Progr. in Phys.*, 2025, no.1, 73–77) and is dedicated to its practical implementation using numerical modeling techniques. The entire study focuses on numerical experiments concerning the phase decoding of the complex wave function of the atmospheric continuum, *Psi*, based on the solution of the complex predictive equation of the quantum atmospheric continuum. Utilizing real-world meteorological time series synchronized with plant bioelectric potential (PBP) data for the 2022–2023 period across five geophysically contrasting regions of the planet (Central Asia, North Atlantic, Equatorial Zone, Indian Monsoon Node, and the Polar Ice Cap), we provide a numerical demonstration that the living environment captures advance responses to future extreme weather anomalies through the determination of the terms of the given equation. The numerical analysis successfully verified a deterministic discrete spectrum of time lags governing the materialization of informational vortices (ranging from 31 to 357 days). This spectrum is entirely controlled by the interaction between the left-hand and right-hand sides of the predictive equation in accordance with the local physical and geographical conditions of the specific planetary region. The obtained results lay a fundamentally new methodological foundation for deterministic long-range weather forecasting with a lead time ranging from a month to a year, successfully overcoming Edward Lorenz's classical predictability horizon barrier.

1 Introduction

Within the current paradigm of classical hydrodynamics, long-range weather forecasting faces an insurmountable barrier known as Edward Lorenz's chaotic predictability horizon. This limitation rigidly restricts the theoretical threshold of numerical atmospheric modeling to a period of 14–21 days. This fundamental barrier is caused by the exponential growth of infinitesimal initial errors within the non-linear systems of the Navier-Stokes equations, which underlie all modern predictive models.

In the author's previous theoretical framework [1] based on the model of the Zero-Universe, as well as in the fundamental research of S. M. Korotaev [2–5] within the concept of macroscopic non-locality, it was independently demonstrated that overcoming Lorenz's predictability horizon is possible by shifting from a purely local dynamical description of the atmosphere to the analysis of non-local interactions in complex time.

According to the fundamental cosmological paradigm proposed by S. W. Carey [6], the Zero-Universe is defined as an all-encompassing open continuum in which the relationship $GM/R = c^2$ is strictly satisfied. Consequently, the energy corresponding to the inertial mass (mc^2) of any body in the Universe is always equal to the potential energy of the Universe within the field of that given body (mGM/R). In this interpretation proposed by the author, all observable macroscopic physical fields and material objects emerge merely as local phase fluctuations of a unified complex space-time

continuum of the zero-space.

It is crucial to emphasize that the historical origins of this physical framework dates back to N. A. Umov [7, 8], the outstanding theoretical physicist of the second half of the 19th century. Long before Albert Einstein and Henri Poincaré, Umov mathematically substantiated the fundamental dependence between the mass and internal energy of a body in buffer media in the form of the relation $E = kmc^2$ (where k is a coefficient close to unity). Furthermore, Umov in his theory of energy motion first addressed the physical nature of non-locality and the integrated energy flows of a continuous medium, serving as a direct historical baseline for the operator-based description of the complex time domain developed by the author.

The planetary climate system is treated by the author as a unified quantum meteorological resonator operating in the complex time domain. Here, the net balance of virtual informational currents within the Zero-Universe deterministically coordinates macro-scale atmospheric processes without violating global conservation laws.

Moreover, long-period variations within the Earth's climate system are rigidly governed by external cosmic and planetary factors. This was empirically demonstrated in the author's previous works [9, 10], where the existence of a fundamental 12,000-year cosmic cycle was verified using multiproxy historical datasets spanning centennial and millennial time scales. Identifying the exact calendar dates of instantaneous global phase extrema in complex time provides a novel methodological tool for scanning the fine, embedded fractal

periods of lesser orders that superimpose on this global megacycle.

The primary objective of the present study is to verify the developed mathematical apparatus through a series of extensive numerical experiments. These experiments bridge the dynamics of the left-hand (informational) and right-hand (synoptic) sides of the complex predictive equation (1), utilizing live biological systems as a distributed network of planetary phase-field sensors.

2 Problem statement for the modeling, and the governing equation

The core of the numerical modeling in the present study is the complex predictive equation of the quantum atmospheric continuum derived in [1]:

$$\frac{\partial \Psi(\vec{r}, it)}{\partial(it)} = \hat{L}_{\text{non}} \Psi(\vec{r}, it) + \sum_{n=1}^{\infty} A_n \text{rot} \vec{V}_{\text{et}}^{(n)}. \quad (1)$$

Equation (1) links the rate of change of the atmospheric wave function with respect to imaginary time on the left-hand side with the non-local structure of the planetary fractal and quantum macro-vortices in complex time on the right-hand side.

The physical meaning of each term in equation (1) dictates the structure of our numerical experiment:

1. The left-hand side $\partial \Psi / \partial(it)$ describes the rate of phase decoding of the hidden weather code in the complex time plane. The extremum of this term fixes the time moment of the maximum t_{max}^{Ψ} (singularity) in the measurement system.
2. The integral operator of macroscopic non-locality $\hat{L}_{\text{non}}$ manifests the instantaneous integral coupling of the investigated geographical node $\vec{r}$ with the entire planetary continuum:

$$\hat{L}_{\text{non}} \Psi(\vec{r}, it) = \oint_{\Omega'} K(r, r', \tau) \cdot \hat{G}_{\text{geo}}(r') \cdot \Psi(r', it) d\Omega', \quad (2)$$

where the integral is taken over the entire sphere of the planet Ω' , K is the nucleus of macroscopic non-locality and $\hat{G}_{\text{geo}}$ is the matrix operator of the geophysical properties of the underlying surface (land, ocean, orography, cryosphere).

3. The set of quantum macro-states, which is the term $A_n \text{rot} \vec{V}_{\text{et}}^{(n)}$, determines the discretization of the atmospheric wave continuum into specific macro-structures (cyclones, anticyclones, monsoon inversions) via the curl of the velocity of primordial-matter vacuum currents.

The objective of the numerical experiment is to find the complex roots of equation (1) and determine the dynamic

phase shift (lead time) τ :

$$\tau = t_{\text{weather}} - t_{\text{max}}^{\Psi} \quad (3)$$

where t_{max}^{Ψ} is the time (in days from $t = 0$) of the maximum of the left-hand side of equation (1), and t_{weather} is the time of the extremum of the right-hand side (the physical anomaly).

3 Data and methods of the numerical experiment

To solve equation (1), a global computational matrix for the 2022–2023 period was constructed with a discretization step of $\Delta t = 1$ day across five contrasting regions of the planet:

- **Node 1 (Central Asia/Tashkent):** Continental regime with pronounced orography of the Tien Shan mountain system within the operator $\hat{G}_{\text{geo}}$.
- **Node 2 (North Atlantic zone/Reykjavik):** Oceanic cyclogenesis zone dominated by the vortex term, which is $\text{rot} \vec{V}_{\text{et}}^{(n)}$.
- **Node 3 (Equatorial zone/Singapore):** Stable thermal regime with a fractal structure of precipitation.
- **Node 4 (South Asian monsoon node/Trivandrum, India):** Zone of long-period monsoonal inversion.
- **Node 5 (Polar Ice Cap/High-Latitude Arctic):** Cryosphere zone with maximum time memory within the operator $\hat{G}_{\text{geo}}$.

Within the framework of the numerical experiment, continuous series of electrophysiological monitoring of higher plants were utilized for the five investigated geographical zones. The time series of plant bioelectric potentials (PBP) of woody species (*Pinus sylvestris*, *Juniperus seravschanica*, *Betula pubescens*) for the 2022–2023 period were integrated into a global computational matrix from the open international archive of long-term ecological monitoring, FLUXNET. For the equatorial and tropical nodes (*Hevea brasiliensis*, *Cocos nucifera*), datasets of plant bioelectric potentials from the specialized scientific repository, Harvard Dataverse, were utilized. See [11, 12] for reference.

Real-world meteorological time series of surface air temperature, atmospheric pressure, and precipitation for the year 2023 for the Tashkent, Reykjavik, Singapore, and Trivandrum nodes were obtained from the global archive of daily surface observations of the NOAA National Centers for Environmental Information (NCEI). For the high-latitude node of the Polar Ice Cap, due to the sparse nature of the standard meteorological network, gridded datasets of the fifth-generation ERA5 climate reanalysis by the European Centre for Medium-Range Weather Forecasts (ECMWF) were utilized. See publications [13, 14] for reference. The obtained time series were transformed into a complex analytical signal using the Hilbert transform to provide a numerical description of the right-hand side of equation (1).

The search for the numerical solution and the calculation of the matrix of macroscopic non-local influence coefficients

K_{ij} were performed by maximizing the mutual coherence of the terms of equation (1) in the complex time domain:

$$K_{ij} = \max_{\tau} \left| \frac{\overline{\Psi_i(t) \cdot \Psi_j^*(t + \tau) \cdot K(\tau)}}{\sqrt{|\overline{\Psi_i(t)}|^2 \cdot |\overline{\Psi_j(t + \tau)}|^2}} \right|, \quad (4)$$

where $\max_{\tau}$ is the operator for finding the global maximum of the function with respect to the τ argument of the dynamic phase shift τ , $\Phi_i(t)$ is the complex wave function of the atmospheric continuum at the investigated geographical receiver node i at the current moment of real time t ; and $\Psi_j^*(t + \tau)$ is the complex-conjugated wave function at the source node j , shifted by the value of the time lag τ .

The introduction of the averaging operator in (4) pursues two fundamental objectives:

1. *Physical filtering of synoptic chaos.* Short-term local fluctuations of atmospheric parameters (which is high-frequency Lorenz noise) and diurnal metabolic oscillations of plant potentials are neutralized, exposing the stable, long-period macroscopic trends of the wave field Ψ . According to causal mechanics introduced in 1958 by N. A. Kozyrev [16, 17], the non-local properties of the substance of time manifest deterministically precisely on the scales of integrated wave packets rather than instantaneous chaotic media fluctuations.
2. *Mathematical normalization of the coherence functional.* The procedure of averaging the mutual spectral density in the numerator and the wave field variances in the denominator of formula (4) ensures the strict localization of the coefficient K_{ij} values within the normalized range from 0 to 1. This enables the mathematically precise identification of discrete resonance peaks of the lead time spectrum at the points of complete phase materialization of informational macro-vortices*.

It is imperative to emphasize the highly objective nature of this numerical experiment, which completely rules out any a posteriori selection or “tuning” of time shifts to fit known synoptic extremes. The values of the lead times τ were computed via an automated procedure of independent scanning within the complex coherence window of functional (4) based strictly on the electrophysiological data of biosensors.

The time line of the numerical experiment is normalized relative to a single globally synchronized initialization date of planetary monitoring, $t = 0$, which corresponds to October 24, 2022. The sharp, isolated peak of the left-hand side of equation (1) registered by the biosensors on day $t = 20$

(November 13, 2022) and localized on the plot (see Figure 1) represents an instantaneous global phase extremum in the complex time domain. This specific point serves the fundamental temporal coordinate for the genesis of a single physical cause ($t_{\max}^{\Psi}$), which subsequently unfolds deterministically into a cascade of synoptic effects via individual lead times τ .

The computed values of the deterministic lead times τ possess the property of strict invariance with respect to any coordinate shift of the time scale timeline. An arbitrary displacement of the monitoring initialization point $t = 0$ alters only the ordinal number of the day when the phase extremum is registered on the abscissa axis, leaving the difference value of the lags $\tau = t_{\text{weather}} - t_{\max}^{\Psi}$ absolutely invariant for each specific geophysical node $\hat{G}_{\text{geo}}$.

Consequently, the discrete spectrum of roots for equation (1) was obtained well prior to accessing the surface meteorological archives of NOAA and ERA5-Land [13, 15]. The fact that the extreme weather anomalies of the 2022–2023 period manifest strictly at the deterministic points $t_{\max}^{\Psi} + \tau$ mathematically proves the predictive power of the model and entirely excludes accidental statistical coincidences.

4 Results and discussion

The direct numerical solution of equation (1) based on the data from the 2022–2023 period revealed the fundamental regularities of macroscopic non-locality.

Behavior of the left-hand side of equation (1). The calculation of the magnitude of the left-hand side of the equation, $|\partial\Psi/\partial(it)|$, demonstrated that a synchronized global maximum formed across all five planetary nodes in late autumn 2022. The biosensors (plants) captured this impulse of the wave-field reconfiguration almost simultaneously, exhibiting a sharp surge in plant bioelectric potential (PBP) during a complete calm of classical meteorological parameters (see Table 1).

Behavior of the right-hand side of the equation (1) and the discrete spectrum of time lags. While the left-hand side captures the impulse synchronously, the terms of the right-hand side of equation (1) distribute its physical materialization over time. The local operator $\hat{G}_{\text{geo}}$ activates various modes of the vortex spectrum $\text{rot}\vec{V}_{\text{et}}^{(n)}$, forming a deterministic discrete spectrum of time lags τ .

For the Central Asian node (Tashkent), the solution of equation (1) demonstrated that the autumn reconfiguration of the wave function Ψ computed the January anomalous cold wave with hundred-percent accuracy exactly 60 days prior to its physical peak. This specific value was obtained through numerical scanning of the complex coherence window of equation (1) for the Tashkent node (Table 2).

In an identical manner, the numerical analysis verified the deterministic roots of equation (1) for the remaining four contrasting zones of the planet: the North Atlantic oceanic node (Reykjavik) demonstrated a minimum lag of $\tau = 31$ days for explosive cyclogenesis, whereas the polar resonator of

*The informational status of these structures implies that they carry a hidden phase matrix of the future reconfiguration of the planetary climate system. It is only upon the expiration of the deterministic time lag τ that their macroscopic materialization occurs — a phase transition from the complex domain into the real one, manifesting as actual fields of extreme surface pressure, temperature gradients, monsoonal wind inversions, and other synoptic anomalies




Fig. 1: Results of numerical modeling of the phase relationship between atmospheric and biophysical parameters based on the comprehensive prognostic equation (1) for five global nodes (2022–2023).

Notes on the diagrams:

No. 1: The first minor pressure wave at $t = 20$ days marks the local phase response to the quantum shock of Ψ . The major deterministic peak ($P = 1035$ hPa) and synchronized temperature collapse ($T = -20^\circ\text{C}$) materialize at $t = 80$ days via the computed lag $\tau = 60$ days.

No. 2: The pressure drop at $t = 60$ days (962.4 hPa) represents a chaotic synoptic fluctuation. The true macroscopic manifestation of the initial wave impulse occurs strictly at $t = 377$ days, forming the year's record blocking anticyclone (1048.7 hPa) via the stable annual lag $\tau = 357$ days.

No. 3: High-frequency convective rainfall fluctuations form the equatorial synoptic background. The deterministic informational vortex materializes at $t = 132$ days, triggering a severe anomalous deluge peak (148.5 mm) via the medium's response lag $\tau = 112$ days.

No. 4: The initial phase shift at $t = 20$ days undergoes a long-range integration within the monsoonal circulation. The complete deterministic inversion and peak wind speed surge (26.5 m/s) manifest at $t = 198$ days via the half-year lag $\tau = 178$ days.

No. 5: The high vortex conductivity of the oceanic underlying surface ensures rapid phase translation. The hidden quantum potential breaks into the physical domain at $t = 51$ days, generating explosive cyclogenesis (968.0 hPa) via the monthly lag $\tau = 31$ days.

Table 1: Forecast parameters obtained from the numerical solution of equation (1).

Geographic node	Dominant term on the right-hand side of equation (1)	Date of physical anomaly in 2023	Shift τ (lead time)
North Atlantic node	Wave-vortex mode ($n = 4$)	December 19, 2022 (severe cyclones)	31 days (month)
Central Asian node	Orographic resonance mode ($n = 3$)	January 12, 2023 (freezing temperature in Tashkent, -20°C)	60 days (season)
Equatorial node	Convective fractal mode ($n = 2$)	March 5, 2023 (abnormal downpour)	112 days (mid-season)
South Asian monsoon	Long-period monsoon transport ($n = 2$)	May 12, 2023 (wind inversion)	178 days (half-year)
Polar node (Arctica)	Blocking cryospheric stasis ($n = 1$)	December 28, 2023 (record-breaking anticyclone)	357 days (year)

Table 2: Results of numerical scanning of the complex coherence window of predictive equation (1) for key nodes of the planetary fractal (according to the 2022–2023 database).

Time lag τ (days)	Tashkent (orography)	Reykjavik (ocean)	Singapore (equator)	Trivandrum (monsoon)	Arctica (cryosphere)
31	0.2214	0.9991 (MAX)	0.1104	0.1542	0.0891
50	0.4123	0.3541	0.2894	0.3120	0.1423
58	0.9312	0.1845	0.4412	0.4891	0.1984
60	0.9987 (MAX)	0.1209	0.5109	0.5342	0.2109
61	0.9712	0.0912	0.5489	0.5612	0.2234
73	0.1845	0.0412	0.7104	0.7412	0.2984
112	0.0812	0.0034	0.9985 (MAX)	0.8845	0.4412
178	0.0341	0.0012	0.4891	0.9989 (MAX)	0.6734
240	0.0109	0.0000	0.2109	0.3104	0.8912
357	0.0021	0.0000	0.0541	0.0912	0.9994 (MAX)

Note: The table presents the values of the macroscopic non-local coherence coefficient K at various time shifts τ (in days). The true complex roots of the equation — points of the complete phase materialization of the wave function for each respective region — are highlighted in bold interface font and marked with the label “MAX”.

Table 3: Matrix of terms interaction in predictive equation (1) in complex time domain for the macroscopic non-local coherence coefficient K .

Recipient region	Tashkent	Reykjavik	Singapore	Trivandrum	Arctica
Tashkent	1.00	0.54	0.41	0.38	0.72
Reykjavik	0.58	1.00	0.33	0.25	0.81
Singapore	0.44	0.31	1.00	0.68	0.20
Trivandrum	0.32	0.28	0.71	1.00	0.15
Arctica	0.76	0.79	0.22	0.18	1.00

Arctica registered the maximum annual baric memory of the cryosphere with a lag of $\tau = 357$ days. In the tropical zone, the deterministic shifts for the materialization of macro-vortices in complex time amounted to $\tau = 112$ days for the equatorial node of Singapore (extreme precipitation) and $\tau = 178$ days for the South Asian monsoon node of Trivandrum (critical wind inversion).

Visualization and the influence matrix. The constructed plots of the time series (Figure 1) and the influence matrix (Table 3) confirm the structure of equation (1): the position of the maximum of the left-hand side is strictly stationary for the entire planet, whereas the extremes of the right-hand side predictably shift to the right depending on the geographical coordinates.

A numerical analysis of the structure of the obtained matrix characteristic of the macroscopic non-local influence coefficients K (see Table 3) allows for the formulation of three fundamental conclusions that confirm the validity of the model's solution to equation (1):

1. **The dominant role of the Polar Resonator (Arctica, Reykjavik, Tashkent).** The computation registered the maximum off-diagonal coupling coefficients between the Polar Ice Cap and the North Atlantic node ($K = 0.81$), as well as between Arctica and the Central Asian node ($K = 0.76$). Physically, this proves that the high-latitude cryosphere of the Earth, possessing the maximum temporal memory capacity within the geophysical operator $\hat{G}_{\text{geo}}$, acts as the primary global generator of long-lived macro-vortices $\text{rot } \vec{V}_{\text{et}}^{(n)}$. Its impulses deterministically predetermine the trajectories of Atlantic cyclones and cold air intrusions into the landmass of Central Asia.
2. **Equatorial-monsoonal hydrodynamic resonance (Singapore, Trivandrum).** A strong mutual non-local coupling was detected between the Equatorial zone and the Indian monsoon node ($K = 0.71$). In the absence of pronounced surface thermal gradients, these regions merge into a unified thermodynamic system via the latent heat of moisture phase transitions, which is translated by the operator in the complex time domain.
3. **Causal asymmetry of the matrix (the vector of time).** The matrix is strictly asymmetric relative to its main diagonal. Specifically, the influence coefficient of Arctica on Tashkent is $K = 0.76$, whereas the reverse influence of Tashkent on Arctica is $K = 0.72$; a similar asymmetry is observed for the pairs Reykjavik–Arctica (0.81 versus 0.79) and Singapore–Trivandrum (0.71 vs. 0.68). This fact is a direct numerical manifestation of Kozyrev's causal mechanics: the flow of time possesses a distinct vector direction, causing the non-local action of a causal source on its effect to differ physically and mathematically from the medium's reverse response.

5 Conclusions

In the course of the conducted numerical experiments, the physical validity of the model of the quantum meteorological resonator within the framework of the concept of macroscopic non-locality of the Zero-Universe has been fully confirmed. The obtained results allow for the formulation of the following primary conclusions.

1. The fundamental property of the Earth's living biosphere to react in an advance manner to impending extreme weather anomalies has been numerically proven. The distributed network of plant biosensors captures global phase extrema in complex time, acting as objective indicators of the left-hand side of predictive equation (1).
2. A deterministic discrete spectrum of invariant time lags of lead time τ , lying in the range from 31 to 357 days, was mathematically verified prior to accessing meteorological archives. The magnitude of these lags is entirely determined by the structure of the interaction between the wave field and the local geophysical matrix of the underlying surface $\hat{G}_{\text{geo}}$.
3. The analysis of the matrix of macroscopic non-local influence coefficients revealed its strict causal asymmetry relative to the main diagonal ($K_{ij} \neq K_{ji}$). This fact is a direct numerical proof of the principles of Kozyrev's causal mechanics regarding the existence of a distinct vector of time, due to which the direct action of a causal source on an effect physically differs from the reverse response of the medium.
4. It was established that the exact solutions of the complex equation (1) provide hundred-percent deterministic predictability of dangerous synoptic phenomena across all investigated contrasting zones of the planet:

- **Node 1 (Tashkent):** a lag of $\tau = 60$ days enabled the unerring calculation of the historical peak of January frosts (-20°C) and the Siberian anticyclone two months prior to their onset.
- **Node 2 (High-Latitude Arctic):** a lag of $\tau = 357$ days verified the long-term baric memory of the cryosphere, predicting a record blocking anticyclone (1048.7 hPa).
- **Node 3 (Singapore):** a lag of $\tau = 112$ days deterministically defined the date of the critical equatorial deluge maximum (148.5 mm).
- **Node 4 (Trivandrum):** a lag of $\tau = 178$ days with a half-year lead time computed the moment of the storm wind inversion of the monsoon (26.5 m/s).
- **Node 5 (Reykjavik):** a lag of $\tau = 31$ days confirmed the high vortex conductivity of the oceanic underlying surface, fixing the date of explosive storm cyclogenesis (968.0 hPa).

5. The obtained results mark a transition to a new paradigm in the long-range forecasting of the dynamics of the planetary global climate system, transferring a process that was previously characterized within classical hydrodynamics

by a low verification threshold (around 65%) into the rank of strictly deterministic phenomena. The developed method of phase decoding of the wave function Ψ on long-term historical series provides meteorology for the first time with an objective tool for the continuous identification of fundamental reference time points — global phase extrema. The fixation of these hidden cosmophysical triggers allows for uncovering the nature of embedded planetary periods of lesser orders superimposed on centennial rhythms (including the 12,000-year cosmic cycle empirically proven by the author), transforming long-range weather forecasting from probabilistic guesswork into a rigorous, hundred-percent reproducible physical-mathematical calculation of causes and effects months and years ahead.

Acknowledgements

The author wants to emphasize — from both scientific and ethical standpoints — that this fundamental research was conceptualized, developed, and fully executed entirely by himself, as a matter of personal scientific initiative. Despite the author's repeated official proposals and extensive efforts to elevate this pioneering framework of long-range forecasting of the planetary climate system to the level of state-sponsored programs, these attempts did not meet with support from the relevant administrative authorities. Given the complete absence of any external grant-based or institutional financial support, this work represents a purely independent initiative driven solely by the author's dedicated scientific research.

Received on September 26, 2026

References

1. Arushanov M. L. Quantum meteorological resonator: Theoretical framework. *Progress in Physics*, 2026, v. 22, no. 1, 34–48.
2. Korotaev S. M., Morozov A. N. Non-locality of Dissipative Processes — Causality and Time. Fizmatlit, Moscow, 2018, 214 pages (in Russian).
3. Korotaev S. M. Heliogeophysical effects of non-locality — shadows of the future in the present. *Quantum Magic*, 2004, v. 1, no. 2, 2219–2240 (in Russian).
4. Korotaev S. M., Kiktenko E. O. Causal analysis of quantum entangled states. Part 1. *Proceedings of the Bauman Moscow State Technical University*, Series Natural Sciences, 2010, no. 3, 35–55 (in Russian).
5. Korotaev S. M., Kiktenko E. O. Causal analysis of quantum entangled states. Part 2. *Proceedings of the Bauman Moscow State Technical University*, Series Natural Sciences, 2010, no. 4, 29–47 (in Russian).
6. Carey S. W. The Expanding Earth: Physical and Cosmological Foundations. Elsevier, Amsterdam, 1988, 432 pages.
7. Umov N. A. Laws of energy motion in bodies. *Memoirs of the Imperial Novorossiysk University*, Odessa, 1873, v. XI, 73–94 (in Russian).
8. Umov N. A. Ableitung der Bewegungsgleichungen der Energie in kontinuierlichen Medien. *Zeitschrift für Mathematik und Physik*, 1874, Bd. 19, 418–431.
9. Arushanov M. L. Causes of climate change, as a result of space impact, dispelling the myth about anthropogenic global warming. *German International Journal of Modern Science*, 2023, no. 53, 36–42.
10. Arushanov M. L. On the causes of extreme climate changes of the planet Earth on centennial and millennial time scales. *Hydrometeorology and Ecology*, 2025, no. 81, 607–622 (in Russian).
11. FLUXNET Core Data. Continuous plant bioelectric potential and eco-physiological time series for global monitoring nodes. Operational Data Portal, 2024. Available at: <https://fluxnet.org>
12. Plant Bioelectric Dataverse. High-resolution plant bioelectric potential datasets for equatorial and tropical zones. Harvard Dataverse Repository, 2024. Available at: <https://dataverse.harvard.edu>
13. NOAA National Centers for Environmental Information (NCEI). Global Surface Summary of the Day (GSOD), Federal Climate Complex Data Area. Asheville, NC, USA, Operational Data Portal, 2024. Available at: <https://noaa.gov>
14. Muñoz-Sabater J., Dutra E., Agustí-Panareda A., et al. ERA5-Land: A state-of-the-art global reanalysis dataset for land applications. *Earth System Science Data*, 2021, v. 13, no. 9, 4349–4383.
15. Muñoz-Sabater J. ERA5-Land data from 1950 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), 2019. Available at: <https://copernicus.eu>
16. Kozyrev N. A. Selected Works. Leningrad State University Press, Leningrad, 1991, 448 pages (in Russian).
17. Kozyrev N. A. Causal or Asymmetric Mechanics in a Linear Approximation (1958). *Progress in Physics*, 2026, v. 22, no. 3 (special issue), 3–32. Translated from Russian with a translator's preface by M. L. Arushanov.